


**Quantili x_p della
distribuzione normale**

$$P = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_p} \exp(-x^2/2) dx$$


P	0	1	2	3	4	5	6	7	8	9
0.0	$-\infty$	-2.33	-2.05	-1.88	-1.75	-1.64	-1.55	-1.48	-1.41	-1.34
0.1	-1.28	-1.23	-1.17	-1.13	-1.08	-1.04	-0.99	-0.95	-0.92	-0.88
0.2	-0.84	-0.81	-0.77	-0.74	-0.71	-0.67	-0.64	-0.61	-0.58	-0.55
0.3	-0.52	-0.50	-0.47	-0.44	-0.41	-0.39	-0.36	-0.33	-0.31	-0.28
0.4	-0.25	-0.23	-0.20	-0.18	-0.15	-0.13	-0.10	-0.08	-0.05	-0.03
0.5	0.00	0.03	0.05	0.08	0.10	0.13	0.15	0.18	0.20	0.23
0.6	0.25	0.28	0.31	0.33	0.36	0.39	0.41	0.44	0.47	0.50
0.7	0.52	0.55	0.58	0.61	0.64	0.67	0.71	0.74	0.77	0.81
0.8	0.84	0.88	0.92	0.95	0.99	1.04	1.08	1.13	1.17	1.23
0.9	1.28	1.34	1.41	1.48	1.55	1.64	1.75	1.88	2.05	2.33

**Valori Percentuali $\chi^2(p)$
per la distribuzione χ^2
con ν gradi di libertà
(area tratteggiata = p)**



ν	.995	.99	.975	.95	.90	.75	.50	.25	.10	.05	.025	.01	.005
1	7.88	6.63	5.02	3.84	2.71	1.32	.455	.102	.0158	.0039	.0010	.0002	.0000
2	10.6	9.21	7.38	5.99	4.61	2.77	1.39	.575	.211	.103	.0506	.0201	.0100
3	12.8	11.3	9.35	7.81	6.25	4.11	2.37	1.21	.584	.352	.216	.115	.0717
4	14.9	13.3	11.1	9.49	7.78	5.39	3.36	1.92	1.06	.711	.484	.297	.207
5	16.7	15.1	12.8	11.1	9.24	6.63	4.35	2.67	1.61	1.15	.831	.554	.412
6	18.5	16.8	14.4	12.6	10.6	7.84	5.35	3.45	2.20	1.64	1.24	.872	.676
7	20.3	18.5	16.0	14.1	12.0	9.04	6.35	4.25	2.83	2.17	1.69	1.24	.989
8	22.0	20.1	17.5	15.5	13.4	10.2	7.34	5.07	3.49	2.73	2.18	1.65	1.34
9	23.6	21.7	19.0	16.9	14.7	11.4	8.34	5.90	4.17	3.33	2.70	2.09	1.73
10	25.2	23.2	20.5	18.3	16.0	12.5	9.34	6.74	4.87	3.94	3.25	2.56	2.16
11	26.8	24.7	21.9	19.7	17.3	13.7	10.3	7.58	5.58	4.57	3.82	3.05	2.60
12	28.3	26.2	23.3	21.0	18.5	14.8	11.3	8.44	6.30	5.23	4.40	3.57	3.07
13	29.8	27.7	24.7	22.4	19.8	16.0	12.3	9.30	7.04	5.89	5.01	4.11	3.57
14	31.3	29.1	26.1	23.7	21.1	17.1	13.3	10.2	7.79	6.57	5.63	4.66	4.07
15	32.8	30.6	27.5	25.0	22.3	18.2	14.3	11.0	8.55	7.26	6.26	5.23	4.60
16	34.3	32.0	28.8	26.3	23.5	19.4	15.3	11.9	9.31	7.96	6.91	5.81	5.14
17	35.7	33.4	30.2	27.6	24.8	20.5	16.3	12.8	10.1	8.67	7.56	6.41	5.70
18	37.2	34.8	31.5	28.9	26.0	21.6	17.3	13.7	10.9	9.39	8.23	7.01	6.26
19	38.6	36.2	32.9	30.1	27.2	22.7	18.3	14.6	11.7	10.1	8.91	7.63	6.84
20	40.0	37.6	34.2	31.4	28.4	23.8	19.3	15.5	12.4	10.9	9.59	8.26	7.43
21	41.4	38.9	35.5	32.7	29.6	24.9	20.3	16.3	13.2	11.6	10.3	8.90	8.03
22	42.8	40.3	36.8	33.9	30.8	26.0	21.3	17.2	14.0	12.3	11.0	9.54	8.64
23	44.2	41.6	38.1	35.2	32.0	27.1	22.3	18.1	14.8	13.1	11.7	10.2	9.26
24	45.6	43.0	39.4	36.4	33.2	28.2	23.3	19.0	15.7	13.8	12.4	10.9	9.89
25	46.9	44.3	40.6	37.7	34.4	29.3	24.3	19.9	16.5	14.6	13.1	11.5	10.5
26	48.3	45.6	41.9	38.9	35.6	30.4	25.3	20.8	17.3	15.4	13.8	12.2	11.2
27	49.6	47.0	43.2	40.1	36.7	31.5	26.3	21.7	18.1	16.2	14.6	12.9	11.8
28	51.0	48.3	44.5	41.3	37.9	32.6	27.3	22.7	18.9	16.9	15.3	13.6	12.5
29	52.3	49.6	45.7	42.6	39.1	33.7	28.3	23.6	19.8	17.7	16.0	14.3	13.1
30	53.7	50.9	47.0	43.8	40.3	34.8	29.3	24.5	20.6	18.5	16.8	15.0	13.8
40	66.8	63.7	59.3	55.8	51.8	45.6	39.3	33.7	29.1	26.5	24.4	22.2	20.7
50	79.5	76.2	71.4	67.5	63.2	56.3	49.3	42.9	37.7	34.8	32.4	29.7	28.0
60	92.0	88.4	83.3	79.1	74.4	67.0	59.3	52.3	46.5	43.2	40.5	37.5	35.5
70	104.2	100.4	95.0	90.5	85.5	77.6	69.3	61.7	55.3	51.7	48.8	45.4	43.3
80	116.3	112.3	106.6	101.9	96.6	88.1	79.3	71.1	64.3	60.4	57.2	53.5	51.2
90	128.3	124.1	118.1	113.1	107.6	98.6	89.3	80.6	73.3	69.1	65.6	61.8	59.2
100	140.2	135.8	129.6	124.3	118.5	109.1	99.3	90.1	82.4	77.9	74.2	70.1	67.3


**Valori percentuali (t_p) per
la distribuzione di Student
con ν gradi di libertà
(area tratteggiata = p)**



ν	.995	.99	.975	.95	.90	.80	.75	.70	.60	.55
1	63.66	31.82	12.71	6.31	3.08	1.376	1.000	.727	.325	.158
2	9.92	6.96	4.30	2.92	1.89	1.061	.816	.617	.289	.142
3	5.84	4.54	3.18	2.35	1.64	.978	.765	.584	.277	.137
4	4.60	3.75	2.78	2.13	1.53	.941	.741	.569	.271	.134
5	4.03	3.36	2.57	2.02	1.48	.920	.727	.559	.267	.132
6	3.71	3.14	2.45	1.94	1.44	.906	.718	.553	.265	.131
7	3.50	3.00	2.36	1.90	1.42	.896	.711	.549	.263	.130
8	3.36	2.90	2.31	1.86	1.40	.889	.706	.546	.262	.130
9	3.25	2.82	2.26	1.83	1.38	.883	.703	.543	.261	.129
10	3.17	2.76	2.23	1.81	1.37	.879	.700	.542	.260	.129
11	3.11	2.72	2.20	1.80	1.36	.876	.697	.540	.260	.129
12	3.06	2.68	2.18	1.78	1.36	.873	.695	.539	.259	.128
13	3.01	2.65	2.16	1.77	1.35	.870	.694	.538	.259	.128
14	2.98	2.62	2.14	1.76	1.34	.868	.692	.537	.258	.128
15	2.95	2.60	2.13	1.75	1.34	.866	.691	.536	.258	.128
16	2.92	2.58	2.12	1.75	1.34	.865	.690	.535	.258	.128
17	2.90	2.57	2.11	1.74	1.33	.863	.689	.534	.257	.128
18	2.88	2.55	2.10	1.73	1.33	.862	.688	.534	.257	.127
19	2.86	2.54	2.09	1.73	1.33	.861	.688	.533	.257	.127
20	2.84	2.53	2.09	1.72	1.32	.860	.687	.533	.257	.127
21	2.83	2.52	2.08	1.72	1.32	.859	.686	.532	.257	.127
22	2.82	2.51	2.07	1.72	1.32	.858	.686	.532	.256	.127
23	2.81	2.50	2.07	1.71	1.32	.858	.685	.532	.256	.127
24	2.80	2.49	2.06	1.71	1.32	.857	.685	.531	.256	.127
25	2.79	2.48	2.06	1.71	1.32	.856	.684	.531	.256	.127
26	2.78	2.48	2.06	1.71	1.32	.856	.684	.531	.256	.127
27	2.77	2.47	2.05	1.70	1.31	.855	.684	.531	.256	.127
28	2.76	2.47	2.05	1.70	1.31	.855	.683	.530	.256	.127
29	2.76	2.46	2.04	1.70	1.31	.854	.683	.530	.256	.127
30	2.75	2.46	2.04	1.70	1.31	.854	.683	.530	.256	.127
40	2.70	2.42	2.02	1.68	1.30	.851	.681	.529	.255	.126
60	2.66	2.39	2.00	1.67	1.30	.848	.679	.527	.254	.126
120	2.62	2.36	1.98	1.66	1.29	.845	.677	.526	.254	.126
∞	2.58	2.33	1.96	1.645	1.28	.842	.674	.524	.253	.126

Distribuzione di FISHER.
Valori dei frattili $F_p(\nu_1, \nu_2)$
 [ν_1, ν_2 gradi di libertà dei
 campioni], dati da:

$$P = \int_0^{F_p} f(F) dF = 0.90$$




ν_2	1	2	3	4	5	6	8	10	12	14	16	20	30	40	50	∞
1	39.86	49.50	53.59	55.83	57.24	58.20	59.44	60.19	60.70	61.07	61.35	61.74	62.26	62.53	62.70	63.33
2	8.53	9.00	9.16	9.24	9.29	9.33	9.37	9.39	9.41	9.42	9.43	9.44	9.46	9.47	9.47	9.49
3	5.54	5.46	5.39	5.34	5.31	5.29	5.25	5.23	5.22	5.20	5.20	5.18	5.17	5.16	5.15	5.13
4	4.54	4.32	4.19	4.11	4.05	4.01	3.96	3.92	3.90	3.88	3.86	3.84	3.82	3.80	3.80	3.76
5	4.06	3.78	3.62	3.52	3.45	3.40	3.34	3.30	3.27	3.25	3.23	3.21	3.17	3.16	3.15	3.10
6	3.78	3.46	3.29	3.18	3.11	3.05	2.98	2.94	2.90	2.88	2.86	2.84	2.80	2.78	2.77	2.72
7	3.59	3.26	3.07	2.96	2.88	2.83	2.75	2.70	2.67	2.64	2.62	2.59	2.56	2.54	2.52	2.47
8	3.46	3.11	2.92	2.81	2.73	2.67	2.59	2.54	2.50	2.48	2.45	2.42	2.38	2.36	2.35	2.29
9	3.36	3.01	2.81	2.69	2.61	2.55	2.47	2.42	2.38	2.35	2.33	2.30	2.25	2.23	2.22	2.16
10	3.29	2.92	2.73	2.61	2.52	2.46	2.38	2.32	2.28	2.26	2.23	2.20	2.16	2.13	2.12	2.06
11	3.23	2.86	2.66	2.54	2.45	2.39	2.30	2.25	2.21	2.18	2.16	2.12	2.08	2.05	2.04	1.97
12	3.18	2.81	2.61	2.48	2.39	2.33	2.24	2.19	2.15	2.12	2.09	2.06	2.01	1.99	1.97	1.90
13	3.14	2.76	2.56	2.43	2.35	2.28	2.20	2.14	2.10	2.07	2.04	2.01	1.96	1.93	1.92	1.85
14	3.10	2.73	2.52	2.39	2.31	2.24	2.15	2.10	2.05	2.02	2.00	1.96	1.91	1.89	1.87	1.80
15	3.07	2.70	2.49	2.36	2.27	2.21	2.12	2.06	2.02	1.99	1.96	1.92	1.87	1.85	1.83	1.76
16	3.05	2.67	2.46	2.33	2.24	2.18	2.09	2.03	1.99	1.95	1.93	1.89	1.84	1.81	1.79	1.72
17	3.03	2.64	2.44	2.31	2.22	2.15	2.06	2.00	1.96	1.93	1.90	1.86	1.81	1.78	1.76	1.69
18	3.01	2.62	2.42	2.29	2.20	2.13	2.04	1.98	1.93	1.90	1.87	1.84	1.78	1.75	1.74	1.66
19	2.99	2.61	2.40	2.27	2.18	2.11	2.02	1.96	1.91	1.88	1.85	1.81	1.76	1.73	1.71	1.63
20	2.91	2.59	2.38	2.25	2.16	2.09	2.00	1.94	1.89	1.86	1.83	1.79	1.74	1.71	1.69	1.61
25	2.92	2.53	2.32	2.18	2.09	2.02	1.93	1.87	1.82	1.79	1.76	1.72	1.66	1.63	1.61	1.52
30	2.88	2.49	2.28	2.14	2.05	1.98	1.88	1.82	1.77	1.74	1.71	1.67	1.61	1.51	1.55	1.46
35	2.85	2.46	2.25	2.11	2.02	1.95	1.85	1.79	1.74	1.70	1.67	1.63	1.57	1.53	1.51	1.41
40	2.84	2.44	2.23	2.09	2.00	1.93	1.83	1.76	1.71	1.68	1.65	1.61	1.54	1.51	1.48	1.38
45	2.82	2.42	2.21	2.07	1.98	1.91	1.81	1.74	1.70	1.66	1.63	1.58	1.52	1.48	1.46	1.35
50	2.81	2.41	2.20	2.06	1.97	1.90	1.80	1.73	1.68	1.64	1.61	1.57	1.50	1.46	1.44	1.33
∞	2.71	2.30	2.08	1.94	1.85	1.77	1.67	1.60	1.55	1.50	1.47	1.42	1.34	1.30	1.26	1.00

I valori $F_{0.05}, F_{0.025}, F_{0.005}$ si possono calcolare tramite la formula:

$$F_{\alpha}(\nu_1, \nu_2) = \frac{1}{F_{1-\alpha}(\nu_2, \nu_1)}$$

Distribuzione di FISHER.
Valori dei frattili $F_p(\nu_1, \nu_2)$
 [ν_1, ν_2 gradi di libertà dei
 campioni], dati da:

$$P = \int_0^{F_p} f(F) dF = 0.95$$




ν_2	1	2	3	4	5	6	8	10	12	14	16	20	30	40	50	∞
1	161.4	199.5	215.6	224.5	230.2	234.0	238.9	241.9	243.9	245.4	246.5	248.0	250.1	251.0	252.0	254.4
2	18.51	19.00	19.16	19.25	19.30	19.33	19.37	19.40	19.41	19.42	19.43	19.44	19.46	19.47	19.47	19.50
3	10.13	9.55	9.28	9.12	9.01	8.94	8.85	8.79	8.75	8.71	8.69	8.66	8.62	8.59	8.58	8.53
4	7.71	6.94	6.59	6.39	6.26	6.16	6.04	5.96	5.91	5.81	5.84	5.80	5.75	5.72	5.70	5.63
5	6.61	5.79	5.41	5.19	5.05	4.95	4.82	4.74	4.68	4.64	4.60	4.56	4.50	4.46	4.44	4.37
6	5.99	5.14	4.76	4.53	4.39	4.28	4.15	4.06	4.00	3.96	3.92	3.87	3.81	3.77	3.75	3.67
7	5.59	4.74	4.35	4.12	3.97	3.87	3.73	3.64	3.57	3.53	3.49	3.44	3.38	3.34	3.32	3.23
8	5.32	4.46	4.07	3.84	3.89	3.58	3.44	3.35	3.28	3.24	3.20	3.15	3.08	3.04	3.02	2.93
9	5.12	4.26	3.86	3.62	3.48	3.37	3.23	3.14	3.07	3.03	2.99	2.94	2.86	2.83	2.80	2.71
10	4.96	4.10	3.71	3.48	3.33	3.22	3.07	2.98	2.91	2.86	2.83	2.77	2.70	2.66	2.64	2.54
11	4.84	3.98	3.59	3.36	3.20	3.09	2.95	2.85	2.79	2.74	2.10	2.65	2.57	2.53	2.51	2.40
12	4.75	3.89	3.49	3.26	3.11	3.00	2.85	2.75	2.69	2.64	2.60	2.54	2.47	2.43	2.40	2.30
13	4.67	3.81	3.41	3.18	2.03	2.92	2.77	2.67	2.60	2.55	2.51	2.46	2.38	2.34	2.31	2.21
14	4.80	3.74	3.34	3.11	2.96	2.85	2.70	2.60	2.53	2.48	2.44	2.39	2.31	2.27	2.24	2.13
15	4.54	3.68	3.29	3.06	2.90	2.79	2.64	2.54	2.48	2.42	2.38	2.33	2.25	2.20	2.18	2.07
16	4.49	3.63	3.24	3.01	2.85	2.74	2.59	2.49	2.42	2.37	2.33	2.28	2.19	2.15	2.12	2.01
17	4.45	3.59	3.20	2.96	2.81	2.70	2.55	2.45	2.38	2.33	2.29	2.23	2.15	2.10	2.08	1.96
18	4.41	3.55	3.16	2.93	2.77	2.66	2.51	2.41	2.34	2.29	2.25	2.19	2.11	2.06	2.04	1.92
19	4.38	3.52	3.13	2.90	2.74	2.63	2.48	2.38	2.31	2.26	2.21	2.16	2.07	2.03	2.00	1.58
20	4.35	3.49	3.10	2.87	2.71	2.60	2.45	2.35	2.28	2.22	2.18	2.12	2.04	1.99	1.97	1.84
25	4.24	3.39	2.99	2.76	2.60	2.49	2.34	2.24	2.16	2.11	2.07	2.01	1.92	1.87	1.84	1.71
30	4.17	3.32	2.92	2.69	2.53	2.42	2.27	2.16	2.09	2.04	1.99	1.93	1.84	1.79	1.76	1.62
35	4.12	3.27	2.87	2.64	2.49	2.37	2.22	2.11	2.04	1.99	1.94	1.88	1.79	1.74	1.70	1.56
40	4.08	3.23	2.84	2.61	2.45	2.34	2.18	2.08	2.00	1.95	1.90	1.84	1.74	1.69	1.66	1.51
45	4.06	3.20	2.81	2.58	2.42	2.31	2.15	2.05	1.97	1.92	1.87	1.81	1.71	1.66	1.63	1.47
50	4.03	3.18	2.79	2.56	2.40	2.29	2.13	2.03	1.95	1.89	1.85	1.78	1.69	1.63	1.60	1.44
∞	3.84	3.00	2.80	2.37	2.21	2.10	1.94	1.83	1.75	1.69	1.64	1.57	1.46	1.39	1.35	1.00

I valori $F_{0.05}, F_{0.025}, F_{0.005}$ si possono calcolare tramite la formula:

$$F_{\alpha}(\nu_1, \nu_2) = \frac{1}{F_{1-\alpha}(\nu_2, \nu_1)}$$

Distribuzione di FISHER.
Valori dei frattili $F_p(\nu_1, \nu_2)$
 [ν_1, ν_2 gradi di libertà dei
 campioni], dati da:

$$P = \int_0^{F_p} f(F) dF = 0.975$$




ν_2	1	2	3	4	5	6	8	10	12	14	16	20	30	40	50	∞
1	647.8	799.4	864.0	399.4	921.6	936.8	956.3	968.3	976.3	982.1	986.5	992.7	1001	1006	1008	1018
2	38.50	39.00	39.17	39.25	39.30	39.33	39.37	39.39	39.41	39.42	39.43	39.44	39.46	39.47	39.48	39.50
3	17.44	16.04	15.44	15.10	14.88	14.73	14.54	14.42	14.34	14.28	14.23	14.17	14.08	14.04	14.01	13.90
4	12.22	10.65	9.98	9.60	9.36	9.20	8.98	8.84	8.75	8.68	8.63	8.56	8.46	8.41	8.38	8.26
5	10.01	8.43	7.76	7.39	7.15	6.98	6.76	6.62	6.52	6.46	6.40	6.33	6.23	6.18	6.14	6.02
6	8.81	7.26	6.60	6.23	5.99	5.82	5.60	5.46	5.37	5.30	5.24	5.17	5.07	5.01	4.98	4.85
7	8.07	6.54	5.89	5.52	5.29	5.12	4.90	4.76	4.67	4.60	4.54	4.47	4.36	4.31	4.28	4.14
8	7.57	6.06	5.42	5.05	4.82	4.65	4.43	4.30	4.20	4.13	4.08	4.00	3.89	3.84	3.81	3.67
9	7.21	5.71	5.08	4.72	4.48	4.32	4.10	3.96	3.87	3.80	3.74	3.67	3.56	3.51	3.47	3.33
10	6.94	5.46	4.83	4.47	4.24	4.07	3.85	3.72	3.62	3.55	3.50	3.42	3.31	3.26	3.22	3.08
11	6.72	5.26	4.63	4.28	4.04	3.88	3.66	3.53	3.43	3.36	3.30	3.23	3.12	3.06	3.03	2.88
12	6.55	5.10	4.47	4.12	3.89	3.73	3.51	3.37	3.28	3.21	3.15	3.07	2.96	2.91	2.87	2.72
13	6.41	4.97	4.35	4.00	3.77	3.60	3.39	3.25	3.15	3.08	3.03	2.95	2.84	2.78	2.74	2.60
14	6.30	4.86	4.24	3.89	3.66	3.50	3.29	3.15	3.05	2.98	2.92	2.84	2.73	2.67	2.64	2.49
15	6.20	4.77	4.15	3.80	3.58	3.41	3.20	3.06	2.96	2.89	2.84	2.76	2.64	2.59	2.55	2.40
16	6.12	4.69	4.08	3.73	3.50	3.34	3.12	2.99	2.89	2.82	2.76	2.68	2.57	2.57	2.47	2.32
17	6.04	4.62	4.01	3.66	3.44	3.26	3.06	2.92	2.82	2.75	2.70	2.62	2.50	2.44	2.41	2.25
18	5.98	4.56	3.95	3.61	3.38	3.22	3.01	2.87	2.77	2.70	2.64	2.56	2.44	2.38	2.35	2.19
19	5.92	4.51	3.90	3.56	3.33	3.17	2.96	2.82	2.72	2.65	2.59	2.51	2.39	2.33	2.30	2.13
20	5.87	4.46	3.86	3.51	3.29	3.13	2.91	2.77	2.68	2.60	2.55	2.46	2.35	2.29	2.25	2.09
25	5.69	4.29	3.69	3.35	3.13	2.97	2.75	2.61	2.51	2.44	2.38	2.30	2.18	2.12	2.08	1.91
30	5.57	4.18	3.59	3.25	3.03	2.81	2.65	2.51	2.41	2.34	2.28	2.20	2.07	2.01	1.97	1.79
35	5.48	4.11	3.52	3.18	2.96	2.80	2.58	2.44	2.34	2.27	2.21	2.12	2.00	1.93	1.89	1.70
40	5.42	4.05	3.46	3.13	2.90	2.74	2.53	2.39	2.29	2.21	2.15	2.07	1.94	1.88	1.83	1.64
45	5.38	4.01	3.42	3.09	2.86	2.70	2.49	2.35	2.25	2.17	2.11	2.03	1.90	1.83	1.79	1.59
50	5.34	3.97	3.39	3.05	2.83	2.67	2.46	2.32	2.22	2.14	2.08	1.99	1.87	1.80	1.75	1.55
∞	5.02	3.69	3.12	2.79	2.57	2.41	2.19	2.05	1.94	1.87	1.80	1.71	1.57	1.48	1.43	1.00

I valori $F_{0.05}, F_{0.025}, F_{0.005}$ si possono calcolare tramite la formula:

$$F_{\alpha}(\nu_1, \nu_2) = \frac{1}{F_{1-\alpha}(\nu_2, \nu_1)}$$

Distribuzione di FISHER.
Valori dei frattili $F_p(\nu_1, \nu_2)$
 [ν_1, ν_2 gradi di libertà dei
 campioni], dati da:

$$P = \int_0^{F_p} f(F) dF = 0.99$$



ν_2	1	2	3	4	5	6	8	10	12	14	16	20	30	40	50	∞
1	4052	5000	5403	5625	5764	5859	5982	6056	6106	6143	6169	6204	6261	6287	6303	6366
2	98.49	99.01	99.18	99.31	99.30	99.33	99.32	99.49	99.41	99.46	99.47	99.43	99.42	99.41	44.59	99.50
3	34.11	30.83	29.46	28.70	28.23	27.90	27.49	27.23	27.05	26.92	26.83	26.69	26.51	26.41	26.30	26.13
4	21.19	17.99	16.69	15.98	15.53	15.21	14.80	14.55	14.38	14.25	14.15	14.02	13.84	13.75	13.69	13.46
5	16.26	13.27	12.06	11.35	10.97	10.67	10.29	10.05	9.89	9.77	9.68	9.55	9.38	9.29	9.24	9.02
6	13.74	10.92	9.78	9.15	8.75	8.47	8.10	7.87	7.72	7.61	7.52	7.40	7.23	7.14	7.09	6.88
7	12.25	9.55	8.45	7.85	7.46	7.19	6.84	6.62	6.47	6.36	6.28	6.16	5.99	5.91	5.86	5.65
8	11.25	8.65	7.59	7.01	6.63	6.37	6.03	5.82	5.67	5.56	5.48	5.36	5.20	5.12	5.07	4.86
9	10.56	8.02	6.99	6.42	6.06	5.80	5.47	5.26	5.11	5.01	4.92	4.81	4.65	4.57	4.52	4.31
10	10.04	7.56	6.55	6.00	5.64	5.39	5.06	4.85	4.71	4.60	4.52	4.41	4.25	4.17	4.12	3.91
11	9.64	7.21	6.22	5.67	5.32	5.07	4.74	4.54	4.40	4.29	4.21	4.10	3.94	3.86	3.81	3.60
12	9.33	6.93	5.95	5.41	5.06	4.82	4.50	4.30	4.16	4.05	3.97	3.86	3.70	3.62	3.57	3.36
13	9.07	6.70	5.74	5.21	4.86	4.62	4.30	4.10	3.96	3.86	3.78	3.66	3.51	3.43	3.38	3.17
14	8.86	6.51	5.56	5.04	4.70	4.46	4.14	3.94	3.80	3.70	3.62	3.51	3.35	3.27	3.22	3.00
15	8.68	6.36	5.42	4.89	4.56	4.32	4.00	3.80	3.67	3.56	3.49	3.37	3.21	3.13	3.08	2.87
16	8.53	6.23	5.29	4.77	4.44	4.20	3.89	3.69	3.55	3.45	3.37	3.26	3.10	3.02	2.97	2.75
17	8.40	6.11	5.19	4.67	4.34	4.10	3.79	3.59	3.46	3.35	3.21	3.16	3.00	2.92	2.87	2.65
18	8.29	6.01	5.09	4.58	4.25	4.01	3.71	3.51	3.37	3.27	3.19	3.08	2.92	2.84	2.78	2.51
19	8.18	5.93	5.01	4.50	4.17	3.94	3.63	3.43	3.30	3.19	3.12	3.00	2.84	2.16	2.71	2.49
20	8.10	5.85	4.94	4.43	4.10	3.87	3.56	3.37	3.23	3.13	3.05	2.94	2.78	2.69	2.64	2.42
25	7.77	5.57	4.68	4.18	3.86	3.63	3.32	3.13	2.99	2.89	2.81	2.10	2.54	2.45	2.40	2.17
30	7.56	5.35	4.51	4.02	3.70	3.47	3.17	2.98	2.84	2.74	2.66	2.55	2.39	2.30	2.25	2.01
35	7.42	5.27	4.40	3.91	3.59	3.37	3.07	2.88	2.74	2.64	2.56	2.44	2.28	2.19	2.14	1.09
40	7.31	5.18	4.31	3.83	3.51	3.29	2.99	2.80	2.66	2.56	2.48	2.37	2.20	2.11	2.06	1.80
45	7.23	5.11	4.25	3.77	3.45	3.23	2.94	2.14	2.61	2.51	2.43	2.31	2.14	2.05	2.00	1.74
50	7.17	5.06	4.20	3.72	3.41	3.19	2.89	2.70	2.56	2.46	2.38	2.27	2.10	2.01	1.95	1.68
∞	6.63	4.61	3.78	3.32	3.02	2.80	2.51	2.32	2.18	2.08	2.00	1.88	1.70	1.59	1.52	1.00

I valori $F_{0.05}, F_{0.025}, F_{0.005}$ si possono calcolare tramite la formula:

$$F_{\alpha}(\nu_1, \nu_2) = \frac{1}{F_{1-\alpha}(\nu_2, \nu_1)}$$